arXiv:1701.00653v2 [math.LO] 4 Mar 2017

ON A QUESTION OF SILVER ABOUT GAP-TWO CARDINAL TRANSFER PRINCIPLES

MOHAMMAD GOLSHANI AND SHAHRAM MOHSENIPOUR

Abstract. Assuming the existence of a Mahlo cardinal, we produce a generic extension of Gödel's constructible universe L, in which the GCH holds and the transfer principles $(\aleph_2, \aleph_0) \to (\aleph_3, \aleph_1)$ and $(\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0)$ fail simultaneously. The result answers a question of Silver from 1971. We also extend our result to higher gaps.

1. Introduction

In this paper we study cardinal transfer principles introduced by Vaught [6], [7], and prove some consistency results related to them.

Assume $\mathcal{L}$ is a first order language which contains a unary predicate U. By a (κ, λ)-model for $\mathcal{L}$, we mean a model $\mathcal{M} = (M, U^{\mathcal{M}}, \dots)$, where $|M| = \kappa$ and $|U^{\mathcal{M}}| = \lambda$, where $U^{\mathcal{M}}$ is the interpretation of U in $\mathcal{M}$. Following Devlin [2], we use the notation

$$(\kappa, \lambda) \to (\kappa', \lambda')$$

to mean the following transfer principle:

> For every countable first order language $\mathcal{L}$ as above, and every first order theory T of $\mathcal{L}$, if T has a (κ, λ)-model, then it has a (κ', λ')-model.

For any natural number $n \geq 1$, by the *gap-n-cardinal transfer principle* we mean the statement

$$\forall \kappa \ \forall \lambda \ (\kappa^{+n}, \kappa) \to (\lambda^{+n}, \lambda).$$

In [5], Silver proved the independence of gap-2-cardinal transfer principle. Starting from an inaccessible cardinal, he was able to produce a model in which the cardinal transfer $(\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0)$ fails. His proof is simply as follows: By a result of Vaught [7], there

The first author's research has been supported by a grant from IPM (No. 95030417).

The research of the second author was in part supported by a grant from IPM (No. 95030403).

exists a sentence ϕ_{KH} in a suitable first order language, such that for any infinite cardinal β,

$$\phi_{KH} \text{ has a } (\beta^{++}, \beta)\text{-model} \iff \text{there exists a } \beta^{+}\text{-Kurepa tree.}$$

Now, starting from an inaccessible cardinal κ, Silver shows that in the generic extension by the Levy collapse $\mathrm{Col}(\aleph_1, < \kappa)$, there are no $\aleph_1$-Kurepa trees. If we start with $V = L$, then in the resulting extension, there are $\aleph_2$-Kurepa trees, and so the transfer principle $(\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0)$ fails in it. Similarly if we force with $\mathrm{Col}(\aleph_2, < \kappa)$, then in the extension there are no $\aleph_2$-Kurepa trees, and we can use it to prove the independence of $(\aleph_2, \aleph_0) \to (\aleph_3, \aleph_1)$. The following question was asked by Silver [5].

Question 1.1. [1] *Is it consistent with GCH that both transfer principles* $(\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0)$ *and* $(\aleph_2, \aleph_0) \to (\aleph_3, \aleph_1)$ *fail simultaneously*?

Remark 1.2. *If we drop the GCH assumption from the question, then one can easily answer the above question. Assume* κ *is an inaccessible cardinals and let* $G * H$ *be* $\mathrm{Col}(\aleph_1, < \kappa) * \underset{\sim}{\mathrm{Add}}(\aleph_0, \kappa)$*-generic over* L*. In the generic extension* $L[G * H]$ *there are no* $\aleph_1$*-Kurepa trees (see Devlin* [3]*) but there exists an* $\aleph_2$*-Kurepa tree, and hence by the remarks above, the transfer principle* $(\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0)$ *fails in* $L[G * H]$.

On the other hand $L[G * H]$ *satisfies "*$2^{\aleph_0} = 2^{\aleph_1} = \kappa = \aleph_2$*". Let* $\mathcal{L} = (U, F)$*, where* U *is a unary predicate symbol and* F *is a binary predicate symbol. let* T *be an* $\mathcal{L}$*-theory which says the following:*

(1) $\forall x, y\ F(x, y) \to U(y)$*. In particular, for each* x*,* F *determines a subset* F_x *of* U*, namely,* $F_x = \{y : F(x, y)\}$.
(2) *For all* $x \neq x', F_x \neq F_y$.

Then T *has an* $(\aleph_2, \aleph_0)$ *model but it does not have an* $(\aleph_3, \aleph_1)$*-model (as otherwise we should have* $2^{\aleph_1} \geq \aleph_3$*). Thus the transfer principle* $(\aleph_2, \aleph_0) \to (\aleph_3, \aleph_1)$ *fails in* $L[G * H]$.

We give an affirmative answer to this question by proving the following theorem:

[1] On page 388 of [5], Silver writes "One can also get a *GCH* model in which $(\aleph_7, \aleph_5) \to (\aleph_3, \aleph_1)$ fails and a *GCH* model which $(\aleph_3, \aleph_1) \to (\aleph_7, \aleph_5)$ fails (though I don't see how to get the $\to$ both ways to fail simultaneously)".

Theorem 1.3. *Assume κ is a Mahlo cardinal. Then there is a generic extension of L, the Gödel's constructible universe, in which the GCH holds and the cardinal transfer principles $(\aleph_2, \aleph_0) \to (\aleph_3, \aleph_1)$ and $(\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0)$ fail.*

Then we prove a general model theoretic fact, and use it to extend the above result to higher gaps:

Theorem 1.4. *Assume κ is a Mahlo cardinal. Then there is a generic extension of L in which the GCH holds and for all $n \geq 2$, the cardinal transfer principles $(\aleph_n, \aleph_0) \to (\aleph_{n+1}, \aleph_1)$ and $(\aleph_{n+1}, \aleph_1) \to (\aleph_n, \aleph_0)$ fail.*

Remark 1.5. *Our proofs can be easily extended to get the following consistency result: assume $\alpha < \beta$ are regular cardinals and assume there exists a Mahlo cardinal above them. Then in a generic extension of L, the GCH holds and both transfer principles $(\alpha^{+n}, \alpha) \to (\beta^{+n}, \beta)$ and $(\beta^{+n}, \beta) \to (\alpha^{+n}, \alpha)$ fail.*

In Section 2 we prove Theorem 1.3 and in Section 3, we prove Theorem 1.4. In the last section, we discuss the same problem for the case of gap-1.

2. Proof of Theorem 1.3

In this section we prove Theorem 1.3.

2.1. **On a result of Jensen.** In this subsection we state a result of Jensen [4] and mention some of its basic properties which are needed. Let $\mathcal{L} = \{\in, A, \mathcal{C}\}$, where A is a unary predicate and $\mathcal{C}$ is a function symbol. Let T_J be the following theory in $\mathcal{L}$:

"$ZFC^- + GCH + A^+$ is the largest cardinal+$\mathcal{C}$ is a $\square_{A^+}$-sequence".

By a (κ, λ)-model of T_J we mean a model $\mathcal{M} = (M, \in^{\mathcal{M}}, A^{\mathcal{M}}, \mathcal{C}^{\mathcal{M}})$ of T_J, where $|M| = \kappa$ and $|A^{\mathcal{M}}| = \lambda$.

Theorem 2.1. *(Jensen [4]) Assume $GCH + \diamondsuit_{\beta^+}$ holds, where β is a regular cardinal, and suppose $\kappa > \beta$ is a Mahlo cardinal. Then there is a forcing notion $\mathbb{P}_{\beta,\kappa}$ such that if K is $\mathbb{P}_{\beta,\kappa}$-generic over V, then the following hold in $V[K]$:*

(a) $V[K] \models$ *"GCH"*.

(b) *The principle* $\diamondsuit^+_{\beta^+}$ *holds.*

(c) *The theory* T_J *does not have any* (β^{++},β)*-model.*

Proof. As requested by the referees, we sketch the proof of the theorem, by providing the forcing construction $\mathbb{P}_{\beta,\kappa}$, and refer to [4] for details. Let G be $\mathrm{Col}(\beta^+,<\kappa)$-generic over V, where

$$\mathrm{Col}(\beta^+,<\kappa)=\{p:\beta^+\times\kappa\to\kappa:|p|\leq\beta \text{ and for all } (\alpha,\lambda)\in\mathrm{dom}(p),\ p(\alpha,\lambda)<\lambda\}$$

is the Levy collapse. The next claim is standard.

Claim 2.2. (a) *The forcing* $\mathrm{Col}(\beta^+,<\kappa)$ *is* β^+*-closed and* κ*-c.c.*

(b) $V[G]\models$ *"*$GCH+\diamondsuit_{\beta^+}$*".*

(c) $V[G]\models$ *"*$\kappa=\beta^{++}$ *and* $\square_{\beta^{++}}$ *fails".*

In [4], the following strengthening of Claim 2.2(c) is proved.

Claim 2.3. *In* $V[G]$*, the theory* T_J *has no* (β^{++},β)*-model*

From now on we work in $V[G]$. Let $\mathcal{S}=\langle S_\alpha:\alpha<\beta^+\rangle$ witness $\diamondsuit_{\beta^+}$. For each $\alpha<\beta^+$ let $d_\alpha:\beta\to\alpha$ be an onto function and set $d=\langle d_\alpha:\alpha<\beta^+\rangle$. For $\alpha<\beta^+$ set

$$M_\alpha=L_{\gamma_\alpha}[\mathcal{S}\restriction\alpha+1,d\restriction\alpha+1],$$

where γ_α is the least ordinal $\gamma>\alpha$ such that $\gamma>\sup_{\nu<\alpha}\gamma_\nu$ and

$$L_\gamma[\mathcal{S}\restriction\alpha+1,d\restriction\alpha+1]\models \text{ "}ZFC^-\text{ "}.$$

Define

$$\mathcal{S}^*=\langle S^*_\alpha:\alpha<\beta^+\rangle,$$

where $S^*_\alpha=P(\alpha)\cap M_\alpha$. We find a generic extension of $V[G]$ in which $\mathcal{S}^*$ is a $\diamondsuit^+_{\beta^+}$-sequence.

Let $A\subseteq\kappa$ be such that $L_\kappa[A]=H(\kappa)$ and define the sequence $\langle\rho_\nu:\nu<\kappa\rangle$ by recursion on ν as follows: ρ_ν is the least ordinal $\rho>\beta^+$ such that

- $\rho>\sup_{\xi<\nu}\rho_\xi$.
- $\langle M_\alpha:\alpha<\beta^+\rangle\in L_\rho[A]$.
- $cf(\rho)=\beta^+$.

- $L_\rho[A] \models$"$ZFC^- + \forall x, |x| \leq \beta^+$.

Set $\tilde{\rho}_\nu = \beta^+ \cup \sup_{\xi<\nu} \rho_\xi$,

$$\mathcal{U}_\nu = \langle L_{\rho_\nu}[A], \in, A \cap \rho_\nu, \langle M_\alpha : \alpha < \beta^+ \rangle \rangle,$$

and for $\nu > 0$ set

$$\tilde{\mathcal{U}}_\nu = \bigcup_{\xi<\nu} \tilde{\mathcal{U}}_\xi = \langle L_{\tilde{\rho}_\nu}[A], \in, A \cap \tilde{\rho}_\nu, \langle M_\alpha : \alpha < \beta^+ \rangle \rangle.$$

Then set

$$f_\nu = \text{the } \mathcal{U}_\nu\text{-least bijection } f : \beta^+ \leftrightarrow \tilde{\rho}_\nu.$$

$$a_\xi = \text{the } \xi\text{-th } a \subseteq \beta^+ \text{ in } L_\kappa[A].$$

$$\tilde{a}_\nu = \{(\xi, \mu) : \xi \in a_{f_\nu(\mu)}\}.$$

We are now ready to define the desired forcing notion, that we denote by $Add(\Diamond^+_{\beta^+})$. First we define the forcing notions $Add(\Diamond^+_{\beta^+})_\nu, \nu < \kappa$, which are the building blocks of the main forcing construction [2].

A condition in $Add(\Diamond^+_{\beta^+})_\nu$ is a subset p of β^+ such that

(1) $p \subseteq \beta^+$ is closed and bounded.

(2) $\alpha \in p \implies \tilde{a}_\nu \cap \alpha \in M_\alpha$.

$Add(\Diamond^+_{\beta^+})_\nu$ is ordered by end extension:

$$p \leq q \iff q = p \cap (\max(p) + 1).$$

Let us now define $Add(\Diamond^+_{\beta^+})$. A condition in $Add(\Diamond^+_{\beta^+})$ is a function p such that

(1) $\text{dom}(p) \subseteq \kappa$ and $|\text{dom}(p)| \leq \beta$.

(2) $\forall \nu \in \text{dom}(p), p(\nu) \in Add(\Diamond^+_{\beta^+})_\nu$.

(3) If $\nu \in \text{dom}(p)$, then

  (a) $f''_\nu[\max(p(\nu))] \subseteq \text{dom}(p)$.

  (b) For each $\xi \in f''_\nu[\max(p(\nu))]$, $\max(p(\xi)) \geq \max(p(\nu))$.

  (c) $\alpha \in p(\nu) \implies \tilde{C}_{p,\nu} \cap \alpha \in M_\alpha$, where

$$\tilde{C}_{p,\nu} = \{(\mu, \xi) \in \max(p(\nu)) \times \max(p(\nu)) : \mu \in p(f_\nu(\xi))\}.$$

[2] In [4], the forcing notion $Add(\Diamond^+_{\beta^+})_\nu$ is denoted by $\mathbb{P}^A_\nu$ and the forcing notion $Add(\Diamond^+_{\beta^+})$ is denoted by $\mathbb{P}^A$

The forcing $Add(\diamondsuit^+_{\beta^+})$ is ordered as follows: $p \leq q$ if and only if

$$\mathrm{dom}(p) \supseteq \mathrm{dom}(q) \text{ and for all } \nu \in \mathrm{dom}(q), p(\nu) \leq_{Add(\diamondsuit^+_{\beta^+})_\nu} q(\nu).$$

Let H be $Add(\diamondsuit^+_{\beta^+})$-generic over $V[G]$. The next claim is proved in [4].

Claim 2.4. (a) *$Add(\diamondsuit^+_{\beta^+})$ is β^+-distributive and $\kappa = \beta^{++}$-c.c.".*

(b) *$V[G * H] \models$"GCH".*

(c) *$\mathcal{S}^*$ witnesses that $\diamondsuit^+_{\beta^+}$ holds in $V[G * H]$.*

(d) *The theory T_J does not have a (β^{++}, β)-model in $V[G * H]$.*

Then $\mathbb{P}_{\beta,\kappa} = \mathrm{Col}(\beta^+, < \kappa) * \underset{\sim}{Add}(\diamondsuit^+_{\beta^+})$ is as required. □

Suppose $K = G * H$ is $\mathbb{P}_{\beta,\kappa}$-generic over V. As $\diamondsuit^+_{\beta^+}$ implies the existence of a β^+-Kurepa tree [2], in $V[K]$, we have β^+-Kurepa trees.

2.2. **Completing the proof of Theorem 1.3.** In this subsection we complete the proof of Theorem 1.3. Thus assume $V = L$ and let κ be a Mahlo cardinal. Let λ be the least inaccessible cardinal. So $\lambda < \kappa$. Let G be $\mathrm{Col}(\aleph_1, < \lambda)$-generic over L. Then:

Lemma 2.5. (a) *$L[G] \models$ "There are no $\aleph_1$-Kurepa trees".*

(b) *$L[G] \models$ "GCH holds".*

(c) *$L[G] \models$ "κ is a Mahlo cardinal".*

Proof. (a) and (b) hold by [5], and (c) is clear, as the forcing $\mathrm{Col}(\aleph_1, < \lambda)$ has size $< \kappa$. □

Let K be $\mathbb{P}^{L[G]}_{\aleph_1,\kappa}$-generic over $L[G]$. We show that $L[G * K]$ is the required model. First note that by Theorem 2.1,

$$L[G * K] \models \text{" there exists an } \aleph_2\text{-Kurepa tree"}.$$

But by Lemma 2.5, $L[G] \models$ "There are no $\aleph_1$-Kurepa trees". On the other hand, $L[G] \models$"$\mathbb{P}_{\aleph_1,\kappa}$ is $\lambda = \aleph_2$-distributive", in particular

$$L[G * K] \models \text{ "There are no } \aleph_1\text{-Kurepa trees"}.$$

It follows that

$$L[G * K] \models \text{ " } (\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0) \text{ fails "}.$$

On the other hand, by Theorem 2.1(b), $L[G*K] \models$"T_J does not have an $(\aleph_3, \aleph_1)$-model". We show that T_J has an $(\aleph_2, \aleph_0)$-model in $L[G*K]$. First note that $\aleph_2^{L[G*K]} = \lambda$, which is inaccessible but not Mahlo in L, so it follows from results of Jensen and Solovay (see [2]) that $\square_{\aleph_1}$ holds in both $L[G]$ and $L[G*K]$. Let $\mathcal{C} = \langle C_\alpha : \alpha < \lambda, \lim(\alpha)\rangle \in L[G]$ witness this. Consider the model

$$\mathcal{M} = (H(\lambda)^{L[G]}, \in, \aleph_0, \mathcal{C}),$$

where $\aleph_0$ is considered as the interpretation of A. Then $\mathcal{M}$ is an $(\aleph_2, \aleph_0)$-model of T. So

$$L[G*K] \models \text{ " } (\aleph_2, \aleph_0) \to (\aleph_3, \aleph_1) \text{ fails " }.$$

The theorem follows.

3. A general model theoretic fact and the proof of Theorem 1.4

In this section we prove a general model theoretic fact, and use it to prove Theorem 1.4.

3.1. **A general model theoretic fact.** In this subsection we prove the following lemma and consider some of its consequences.

Lemma 3.1. *Assume $n \geq 1$, $\mathcal{L}$ is a first order language which contains a unary predicate U, and T is a theory in $\mathcal{L}$. Then there are $\mathcal{L}^+ \supseteq \mathcal{L}$ and a theory T^+ in $\mathcal{L}^+$, such that for all infinite cardinals β:*

$$T \text{ has a } (\beta^{+n}, \beta)\text{-model} \iff T^+ \text{ has a } (\beta^{+n+1}, \beta)\text{-model.}$$

Proof. Let $\mathcal{L}^+ = \mathcal{L} \cup \{<, W_0, \ldots, W_n, F_{-1}, F_0, \ldots, F_n\}$ where $<$ is a binary predicate symbol, W_i's are unary predicate symbols, F_{-1} is a binary predicate symbol and F_i's, $0 \leq i \leq n$, are ternary predicate symbols. Let T^+ consists of the following axioms:

(1) ϕ^{W_n}, for each $\phi \in T$, where ϕ^{W_n} is the relativization of ϕ to W_n.

(2) $<$ is a linear ordering of the universe.

(3) Under $<$, each W_i is an initial segment of $W_{i+1}, i < n$, and W_n is an initial segment of the universe (in particular $W_0 \subseteq W_1 \subseteq \cdots \subseteq W_n$).

(4) $U \subseteq W_n$ (i.e., $\forall x(U(x) \to W_n(x))$).

(5) $F_{-1} \subseteq U \times W_0$ defines a bijection from U onto W_0.

(6) For each $0 \leq i < n, F_i \subseteq (W_{i+1} \setminus W_i) \times W_i \times W_{i+1}$ is such that if $x \in W_{i+1} \setminus W_i$, then $\{(y,z) : F_i(x,y,z)\}$ is a bijection from W_i onto $\{z \in W_{i+1} : z < x\}$.

(7) F_n is such that if $x \notin W_n$, then $\{(y,z) : F_n(x,y,z)\}$ is a bijection from W_n onto $\{z : z < x\}$.

Now suppose that T has a (β^{+n}, β)-model $\mathcal{M} = (\beta^{+n}, U^{\mathcal{M}}, \dots)$. Consider the model

$$\mathcal{M}^+ = (\beta^{+n+1}, \mathcal{M}, <, \beta, \dots, \beta^{+n}, f_{-1}, f_0, \dots, f_n),$$

where $f_{-1} : U^{\mathcal{M}} \leftrightarrow \beta$, each $f_i, 0 \leq i \leq n$ is such that for each $\beta^{+i} \leq \gamma < \beta^{+i+1}, \{(\zeta,\eta) : (\gamma,\zeta,\eta) \in f_i\}$ defines a bijection $\beta^{+i} \leftrightarrow \gamma$. It is easily seen that $\mathcal{M}^+$ is a (β^{+n+1}, β)-model for T^+.

Conversely assume that $\mathcal{M}^+$ is a (β^{+n+1}, β)-model for T^+. Consider the model $\mathcal{M}$ which is obtained from $\mathcal{M}^+ \restriction \mathcal{L}$, by replacing its universe with $W_n^{\mathcal{M}^+}$. It follows from (1) that $\mathcal{M}$ is a model of T. We show that it is a (β^{+n}, β)-model. We have $U^{\mathcal{M}} = U^{\mathcal{M}^+}$, which has size β. On the other hand, axioms (4)-(6) can be used to show that $|W_0^{\mathcal{M}^+}| = \beta$, $|W_{i+1}^{\mathcal{M}^+}| \leq |W_i^{\mathcal{M}^+}|^+$ and $|W_m^{\mathcal{M}^+}| \geq \beta^{+n}$, so by induction on $i \leq n$, we have $|W_i^{\mathcal{M}^+}| = \beta^{+i}$. In particular $|W_n^{\mathcal{M}^+}| = \beta^{+n}$, and the result follows. □

Corollary 3.2. *For each $n \geq 1$, the gap-$(n+1)$-cardinal transfer principle implies the gap-n-cardinal transfer principle.*

Remark 3.3. *In personal communication, Ali Enayat informed us that Corollary 3.2 is an immediate consequence of the downward Löwenheim-Skolem theorem, i.e., the fact that if $\mathcal{M} = (M, \dots)$ is an infinite structure in a countable language and X is any subset of M, then there is an elementary substructure $\mathcal{M}_0 = (M_0, \dots)$ of $\mathcal{M}$ that includes X and whose cardinality is $\max\{\aleph_0, |X|\}$. Using this theorem, it is easy to see that every model $\mathcal{M}$ that exhibits a gap-m model, say (κ^{+m}, κ), for some $m > 0$ has an elementary sub-model $\mathcal{M}_0$ that exhibits a gap-n model (κ^{+n}, κ) for all $n < m$.*

3.2. **Proof of Theorem 1.4.** In this subsection we complete the proof of Theorem 1.4. Let $L[G * H]$ be the model obtained in Subsection 2.2. So in $L[G * H]$ both transfer principles $(\aleph_3, \aleph_1) \to (\aleph_2, \aleph_0)$ and $(\aleph_2, \aleph_0) \to (\aleph_3, \aleph_1)$ fail. So, by induction, and using Lemma 3.1, for

each $n \geq 2$, the transfer principles

$$(\aleph_n, \aleph_0) \to (\aleph_{n+1}, \aleph_1)$$

and

$$(\aleph_{n+1}, \aleph_1) \to (\aleph_n, \aleph_0)$$

fail in $L[G * H]$.

4. The case of gap-1 and some problems

In general, we can not hope to prove a result as above for gap-1-cardinal transfer principles. This is because of Vaught's theorem [7] that the transfer principle $(\beta^+, \beta) \to (\aleph_1, \aleph_0)$ is a theorem of ZFC. However we do not know the answer to the following question:

Question 4.1. *Is it consistent that both transfer principles* $(\aleph_2, \aleph_1) \to (\aleph_3, \aleph_2)$ *and* $(\aleph_3, \aleph_2) \to (\aleph_2, \aleph_1)$ *fail simultaneously.*

As we showed in Corollary 3.2, the gap-$(n+1)$-cardinal transfer principle implies the gap-n-cardinal transfer principle.

On the other hand if $L[G]$ is a generic extension of L by the Levy collapse of an inaccessible cardinal κ to $\aleph_2$, then it follows from results of Vaught [7], Chang [1] and Jensen [2] that the gap-1-cardinal transfer principle holds in $L[G]$, while by Silver's result stated in the introduction, the gap-2-cardinal transfer principle fails in $L[G]$. We do not know the answer for higher gaps.

Question 4.2. *Assume* $n > 1$. *Is it consistent that the gap-*n*-cardinal transfer principle holds while the gap-*$(n+1)$*-cardinal transfer principle fails*?

Acknowledgements. The authors would like to thank the referees of the paper for their very helpful comments and corrections. They also thank Ali Enayat for his interest in this work and his Remark 3.3.

References

[1] Chang, C. C. A note on the two cardinal problem. Proc. Amer. Math. Soc. 16 1965 1148-1155.

[2] Devlin, Keith J., Aspects of constructibility. Lecture Notes in Mathematics, Vol. 354. Springer-Verlag, Berlin-New York, 1973. xii+240 pp.

[3] Devlin, Keith J. Kurepa's hypothesis and the continuum. Fund. Math. 89 (1975), no. 1, 23-31.

[4] Jensen, Ronald; Remarks on the Two Cardinal Problem, unpublished note, available at "https://www.mathematik.hu-berlin.de/ raesch/org/jensen.html".

[5] Silver, Jack; The independence of Kurepa's conjecture and two-cardinal conjectures in model theory. 1971 Axiomatic Set Theory (Proc. Sympos. Pure Math., Vol. XIII, Part I, Univ. California, Los Angeles, Calif., 1967) pp. 383-390 Amer. Math. Soc., Providence, R.I.

[6] Vaught, R. L. A Löwenheim-Skolem theorem for cardinals far apart. 1965 Theory of Models (Proc. 1963 Internat. Sympos. Berkeley) pp. 390-401 North-Holland, Amsterdam.

[7] Vaught, R. L. The Löwenheim-Skolem theorem. 1965 Logic, Methodology and Philos. Sci. (Proc. 1964 Internat. Congr.) pp. 81-89 North-Holland, Amsterdam.

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran-Iran.

E-mail address: golshani.m@gmail.com

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran-Iran.

E-mail address: sh.mohsenipour@gmail.com